

Westside Medical Associates, LLP
Ira Daniel Breite, MD
228 W. 82nd Street
New York, NY 10024
(212) 362-6468
Fax: (212) 362-0851
ibreite@westsidedoctors.com

SUPREP INSTRUCTIONS

On the day *before* your procedure...

- You may have a light breakfast or have clear liquids ONLY; please have nothing for dinner.
- DO NOT drink milk
- DO NOT eat or drink anything colored red or purple
- DO NOT drink alcoholic beverages

Any of the following liquids are OK to drink



Water



Strained fruit juices (without pulp) including apple, orange, white grape, or white cranberry



Limeade or lemonade



Coffee or tea. DO NOT use any dairy or non-dairy creamer



Chicken broth



Gelatin desserts without added fruit or toppings (NO RED OR PURPLE)

Click below for more information:

- [Patient Instructions for Use Booklet](#)
- [Instruction Sheet](#)

When to take SUPREP BOWEL PREP KIT

SUPREP Bowel Prep Kit is taken as a split dose (2-day) regimen. You take the first 6-ounce bottle of SUPREP the evening before your colonoscopy and the second 6-ounce bottle of SUPREP the morning of your colonoscopy. It is important to drink the additional water as recommended in the Instructions for Use. Both 6-ounce bottles are required for a complete prep. Your doctor will tell you when it is time to begin to take each dose.

How to take SUPREP BOWEL PREP KIT

- **On the evening before your procedure** (Usually at 7 PM, although for procedures later in the day its up to you), complete steps 1 through 4 using one (1) 6-ounce bottle before going to bed
- **On the morning of your procedure**, repeat steps 1 through 4 using the other 6-ounce bottle. DO NOT HAVE ANYTHING TO DRINK AFTER DOING THIS. MAKE SURE YOU

COMPLETE THE PREP 4 HOURS PRIOR TO THE PROCEDURE. DRINK PLENTY OF FLUIDS (AT LEAST WHAT THE INSTRUCTIONS SAY) PRIOR TO THIS

Both 6-ounce bottles are required for a complete preparation.

IT IS IMPORTANT TO FOLLOW ALL THE STEPS BELOW COMPLETELY.

Step 1:



Pour **ONE** (1) 6-ounce bottle of SUPREP liquid into the mixing container.

Step 2:



Add cool drinking water to the 16-ounce line on the container and mix.

NOTE: Be sure to dilute SUPREP as shown at left before you drink it.

Step 3:



Drink **ALL** the liquid in the container.

Step 4:



You **must** drink two (2) more 16-ounce containers of water over the next 1 hour.

NOTE: You **must** finish drinking the final glass of water at least 1 hour, or as directed, before your procedure.

[Click here for full Prescribing Information and Medication Guide](#)